


Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

# Drug Metabolism and Pharmacokinetics

journal homepage: <http://www.journals.elsevier.com/drug-metabolism-and-pharmacokinetics>

## Short communication

# Novel multiple assessment of hepatocellular drug disposition in a single packaged procedure

Ryosuke Takahashi<sup>a</sup>, Hisashi Ichikawa<sup>a</sup>, Katsuhiko Kanda<sup>b,\*</sup><sup>a</sup> Center for Technology Innovation – Healthcare, Hitachi, Ltd., 2520 Hatoyama, Saitama 350-0395, Japan<sup>b</sup> Science Systems Design Div., Hitachi High-Technologies Corporation, 882, Ichige, Hitachinaka, Ibaraki 312-8504, Japan

## ARTICLE INFO

### Article history:

Received 11 August 2015  
Received in revised form  
10 December 2015  
Accepted 12 December 2015  
Available online 20 December 2015

### Keywords:

Drug disposition  
Hepatocyte  
Uptake  
Basolateral efflux  
Biliary excretion  
Sandwich culture

## ABSTRACT

Better prediction of drug disposition prior to the clinical trial is critical for the efficient development of new drugs. The purpose of this study is to develop a novel multiple assessment methodology of hepatocellular drug disposition from drug uptake to efflux including biliary and basolateral excretion, in a single packaged procedure. We started a sandwich culture using rat primary hepatocytes. After five days culture, the hepatocytes were incubated with a dosing solution including CDF or Rhodamine 123. Three distinct sequences were then performed in parallel: disrupting and maintaining the tight junctions comprising a bile canalicular network at 37 °C, and maintaining the network at 4 °C. Supernatant fractions were collected from each sequence, and followed by the cell lysate collection. The disposition rates of basolateral efflux by diffusion, by transporter-mediation, biliary excretion, and residual cellular fraction of CDF and Rhodamine 123 were 38.2% and 11.0%, 26.6% and 12.1%, 18.6% and 4.9%, and, 16.7% and 72.0%, respectively. CDF was likely to excrete extracellularly whereas Rhodamine 123 tended to remain intracellularly. CDF showed a relatively higher biliary excretion rate than Rhodamine 123. This novel protocol may contribute to improve the predictability of pharmacokinetics eventually in human, and streamline new drug development.

**Chemical compounds:** 5(6)-Carboxy-2',7'-dichlorofluorescein (PubChem CID: 132525); Rhodamine 123 (PubChem CID: 65217).

Copyright © 2015 The Authors. Published by Elsevier Ltd on behalf of The Japanese Society for the Study of Xenobiotics. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Clinically relevant prediction of drug disposition during the early stages of drug development is critical for the efficient development of new drugs. In recent years, two of the primary factors behind the current low clinical success rate of drug candidates (~11%) [13] are lack of efficacy (56%) and safety issues (28%) [2], which have been first detected in the late clinical study stages of the drug development process. In order to reduce the attrition rate, there is a growing need for an in vitro methodology that can clarify the entire pharmacokinetics of a drug eventually applied for better prediction in humans.

Primary hepatocytes have been widely used as effective cell sources for DMPK-related or toxicological study. The sandwich culture system was proposed in the late 1980s as a way to mimic

the hepatocytes of an in vivo environment [3,4]. The sandwich-cultured hepatocytes exhibited the structural polarity with bile canalicular network and the localization of proper hepatobiliary transporters on the network, maintaining liver-specific functions for several days [9,18]. This culture method has been utilized for investigating the hepatic disposition of drugs. In vitro biliary clearance determined in sandwich-cultured hepatocytes correlates well with in vivo biliary clearance [7], and biliary clearance calculation based on the intracellular concentration of a drug has the potential to reflect more the in vivo process than that based on incubation medium [11]. In another report, basolateral and biliary clearance was revealed to be an alternative elimination route in sandwich-cultured hepatocytes by combining biliary excretion and basolateral efflux experiment individually [14].

The objective of this study is to consider these previous reports as a basis and develop an in vitro multiple assessment methodology using hepatocytes to determine the entire picture of drug dispositions including uptake, biliary excretion, and basolateral efflux disposition in a single packaged procedure. The evaluation of these endpoints at the same time in a single procedure has not been

\* Corresponding author.

E-mail addresses: [ryosuke.takahashi.jr@hitachi.com](mailto:ryosuke.takahashi.jr@hitachi.com) (R. Takahashi), [katsuhiko.kanda.pv@hitachi-hightech.com](mailto:katsuhiko.kanda.pv@hitachi-hightech.com) (K. Kanda).

reported. Furthermore, diffusion and transporter-mediated efflux can be distinguished by controlling the specific experimental conditions.

## 2. Materials and methods

### 2.1. Chemicals

5(6)-Carboxy-2,7'-dichlorofluorescein (CDF) and Aprotinin were purchased from Wako Pure Chemical Industries, Ltd. (Osaka, Japan). Rhodamine 123 (Rh 123), Insulin, and Epidermal Growth Factor (EGF) were purchased from Sigma–Aldrich (St. Louis, MO). Penicillin–streptomycin solution and Dexamethasone was purchased from Nakalai Tesque, Inc. (Kyoto, Japan). Matrigel was purchased from BD Bioscience (San Jose, CA). All other chemicals and reagents were of analytical grade and readily available from commercial sources.

### 2.2. Hepatocyte culture

Hepatocytes were isolated from six- to eight-week-old specific viral-pathogen-free male SD rats weighing about 150–250 g (Charles River Japan Inc., Japan) by modified two-step *in situ* collagenase perfusion and purified by Percoll gradient separation [8,15]. Hepatocytes with over 90% viability were used for the following culture.

The hepatocytes were resuspended in William's E medium (Sigma–Aldrich) containing 10% fetal bovine serum, supplemented with 8.6 nM insulin, 255 nM dexamethasone, 50 ng/ml EGF, and 5 KIU/ml aprotinin, and seeded at a density of  $1 \times 10^5$  cells/cm<sup>2</sup> into a type I collagen-coated 24-well culture dish (AGC Techno Glass Co., Ltd., Chiba, Japan). The seeded hepatocytes were incubated in a humidified chamber with 5% CO<sub>2</sub> at 37 °C. After 24 h of post-seeding, the culture medium was replaced with serum-free William's E medium containing Matrigel with the same supplements described above. Subsequently, the culture medium without Matrigel was changed daily. Hepatocytes were cultured for a total of four days.

### 2.3. Fluorescent and phasecontrast microscopy

Sandwich-cultured hepatocytes were rinsed with standard (Std) Hanks' balanced salt solution (HBSS) containing Ca<sup>2+</sup> and Mg<sup>2+</sup> on day 5 and then incubated with a dosing solution including 10 μM CDF or 10 μM Rh 123 in the 5% CO<sub>2</sub> incubator. After 30-min incubation, HBSS was removed and the hepatocytes were washed with Std HBSS. Fluorescent and phasecontrast micrographs of the hepatocytes were acquired with a Zeiss Axiovert 200 inverted optical microscope (Carl Zeiss Microscopy GmbH, Germany).

### 2.4. Disposition study

Sandwich-cultured hepatocytes were rinsed and preincubated in the 37 °C, 5% CO<sub>2</sub> incubator with Std HBSS containing Ca<sup>2+</sup> and Mg<sup>2+</sup> (Std HBSS) on day 5, which the MRP2 expression still maintained in a high rate [16]. Hepatocytes were then incubated with a dosing solution including 10 μM CDF or 10 μM Rh 123 (Fig. 1) in the CO<sub>2</sub> incubator for 30 min. After incubation, the dosing solution was aspirated from the cells and the uptake process was stopped by washing the cells with ice-cold Std HBSS. Subsequently, three different sequences were performed in parallel: one, disrupting the tight junctions comprising a bile canalicular network at 37 °C by Ca<sup>2+</sup>, Mg<sup>2+</sup>-free HBSS with EGTA (seq. 1), two, maintaining the network at 37 °C by Std HBSS (seq. 2), and three, maintaining the network at 4 °C by Std HBSS (seq. 3). The supernatant fractions

were collected from each sequence after 30 min incubation. Finally, cells were lysed with 1% Triton X-100 in H<sub>2</sub>O and cell lysate was collected. The samples were analyzed for relative luminescence unit (RLU) of CDF and Rh 123 by microplate reader.

### 2.5. Data analysis

The disposition rates of each fraction were calculated using the obtained fluorescent data. B2 corresponds to the amount of total basolateral efflux, B3 corresponds to that of basolateral efflux by diffusion, B2 – B3 corresponds to that of basolateral efflux mediated by transporters, B1 – B2 corresponds to that of biliary efflux, and C1 corresponds to that of intracellular residue. In this analysis, B1 + C1 which represents putative intracellular drug amount in the beginning of the efflux fractionation process was counted as 100%. The disposition rate of total basolateral efflux (BEfx-Total), basolateral efflux by diffusion (BEfx-Dif), basolateral efflux by transporter (BEfx-TP), biliary efflux (BCEfx), and intracellular residue (Cell) were calculated by Eqs. (1)–(5), respectively.

$$\text{BEfx-Total} = \text{B2}/(\text{B1} + \text{C1}) \quad (1)$$

$$\text{BEfx-Dif} = \text{B3}/(\text{B1} + \text{C1}) \quad (2)$$

$$\text{BEfx-TP} = (\text{B2} - \text{B3})/(\text{B1} + \text{C1}) \quad (3)$$

$$\text{BCEfx} = (\text{B1} - \text{B2})/(\text{B1} + \text{C1}) \quad (4)$$







$$\text{Cell} = \text{C1}/(\text{B1} + \text{C1}) \quad (5)$$

## 3. Results and discussion

CDF and Rh 123 were independently exposed to sandwich-cultured rat hepatocytes to examine the localization of these two compounds (Fig. 1), which are not subject to hepatic metabolism and primarily excreted in an unchanged form [1,21]. CDF has been used as a model fluorescent compound to evaluate the excretion into bile canaliculi by Multidrug Resistant-associated Protein (MRP) 2 [10,21]. Rh 123, also a fluorescent substance, has been used to determine mitochondrial membrane potential [20] and is currently used to label mitochondria in living cells [5]. CDF was accumulated in the bile canalicular network of sandwich-cultured hepatocytes for 30-min incubation (Fig. 1). In contrast, Rh 123 was localized in the cytoplasm (Fig. 1). These results indicate the characteristics of the two tested compounds.

Multiple drug disposition assay was performed on day 5 according to the protocol described in Materials and methods and Fig. 2. We incubated with CDF or Rh 123 for 30 min at the uptake process. Due to the existing reports, uptake time is generally applied 10–20 min [6,12,19]. Whereas their uptake condition requires two different SCRH buffers with Ca<sup>2+</sup>/Mg<sup>2+</sup> and without Ca<sup>2+</sup>/Mg<sup>2+</sup>, our protocol is performed only with Ca<sup>2+</sup>/Mg<sup>2+</sup>, which is favorable to maintain hepato-cellularly stable condition with intact intercellular adhesion. Accordingly, our uptake condition enables prolonged absorption of the drugs in SCRH. We applied 30 min for efflux experiment because of the results based on the preliminary experiment (data not shown). The time-course of supernatant fluorescence of efflux fractions saturated at about 30 min. We concluded 30 min was suitable for efflux study.

In order to calculate the disposition rate of biliary excretion, the supernatant fraction of B2 was subtracted from the B1 fraction (Fig. 2). The calculation of transporter-mediated basolateral efflux was performed by subtracting B3, a supernatant of the 4 °C

Name	5(6)-Carboxy-2',7' -dichlorofluorescein (CDF)	Rhodamine 123 (Rh 123)
Structure		
Property	MRP2-mediated biliary excretion	Mitochondria membrane accumulation
Phasecontrast micrograph		
Fluorescent micrograph		
	Ex. 484 nm, Em. 519 nm	Ex. 484 nm, Em. 530 nm


**Fig. 1.** Chemical structure, property and micrographic images of 5(6)-carboxy-2,7'-dichlorofluorescein (CDF) and Rhodamine 123. Fluorescent and phasecontrast micrographs of sandwich-cultured rat hepatocytes. Sandwich-cultured hepatocytes were incubated with 10  $\mu$ M CDF (A, B) or with 10  $\mu$ M Rh 123 (C, D) for 30 min. Dosing solution was rinsed with Std HBSS, fluorescent micrograph (A, C) and phasecontrast micrograph (B, D) were taken on day 5.

condition, from the B2 fraction. Thus, the disposition rates of total basolateral efflux (BEfx-Total), basolateral efflux by diffusion (BEfx-Dif), by transporter-mediation (BEfx-TP), biliary excretion (BCEfx), and residual cellular fraction (Cell) of CDF and Rh 123 were  $64.75 \pm 7.39\%$  and  $47.98 \pm 1.61\%$ ,  $38.15 \pm 4.01\%$  and  $10.94 \pm 0.51\%$ ,  $26.60 \pm 7.54\%$  and  $12.11 \pm 1.26\%$ ,  $18.58 \pm 4.60\%$  and  $4.92 \pm 0.82\%$ , and  $16.67 \pm 3.34\%$  and  $72.02 \pm 2.82\%$  according to Eqs. (1)–(5), respectively (Table 1). CDF and Rh 123 are extensively known to be substrates of multiple transporters such as MRP2 [16], MRP3 and P-gp [17], which expressed in sandwich-cultured rat hepatocytes at day 5. CDF was likely to excrete extracellularly whereas Rh 123 tended to remain intracellularly. Additionally, CDF showed a relatively higher biliary excretion rate than Rh 123. These results show the disposition rate of CDF is different from that of Rh 123, which reflects the properties and the localization depending on the compounds illustrated in Fig. 1. We introduced sequence 3 to distinguish the transporter-mediated and passive-diffusion efflux process operated under 4 °C. In order to seek the reliability and

precision of the methodology, the transporter inhibition study using several statins and inhibitors is under evaluation.

The amount of total uptake can be calculated by the sum of the fractionation processes S, B and C in the developed protocol. However, we defined the fractionation process S (Fig. 2) as the process for transition to an apparent steady state of mass transfer, to ensure the following efflux more stable, because our purpose in this experiment is to evaluate the disposition rate based on efflux fractions. Therefore, fraction S was not included as a part of the denominator that corresponds to the total amount of accumulated compounds, and instead, the sum of the following recovered fractions B and C was defined as the total amount, which represents the intracellular total drug amount in the beginning of the efflux study. From this viewpoint, the disposition rates were calculated using Eqs. (1)–(5).

In conclusion, we have developed a multiple assessment methodology for hepatocellular drug disposition. Our methodology can evaluate overall disposition including in vitro uptake, biliary excretion and basolateral efflux, in a single packaged procedure.



**Fig. 2.** Scheme depicting multiple assay protocol for hepatocellular drug disposition. Three distinct sequences were performed in fractionation process B in parallel: one, disrupting tight junctions comprising a bile canalicular network at 37 °C (seq. 1), two, maintaining the network at 37 °C (seq. 2), and three, maintaining the network at 4 °C (seq. 3). Basolateral efflux by diffusion (a) is described by B3, basolateral efflux by transporter (b) is calculated by B2 – B3, biliary excretion (c) is calculated by B1 – B, and intracellular fraction (d) is indicated by C1.

**Table 1**  
Disposition rates of CDF and Rhodamine 123. The disposition rate of basolateral efflux by diffusion (BEfx-Dif), basolateral efflux by transporter (BEfx-TP), biliary efflux (BCEfx), and intracellular residue (Cell) were calculated by Eqs. (1)–(5), respectively. Data represents mean  $\pm$  SD of four individual assays.

Compound	BEfx-Total	BEfx-Dif	BEfx-TP	BCEfx	Cell
%					
CDF	64.75 $\pm$ 7.39	38.15 $\pm$ 4.01	26.60 $\pm$ 7.54	18.58 $\pm$ 4.60	16.67 $\pm$ 3.34
Rhodamine 123	47.98 $\pm$ 1.61	10.95 $\pm$ 0.52	12.11 $\pm$ 1.26	4.92 $\pm$ 0.82	72.02 $\pm$ 2.82
Equation	(1)	(2)	(3)	(4)	(5)

Moreover, the protocol also has a potential to distinguish diffusion and transporter-mediated efflux. In addition, all the compounds are dosed under an identical cellular condition which the bile canalicular network of sandwich-cultured hepatocytes is maintained. The fundamental advantage of this protocol would be expected to allow the processes from the drug uptake to the fractionation process S thoroughly treated under a stable condition for sandwich-cultured hepatocytes, which contributes to precise analyses. Our novel protocol with the advantages above has a potential to contribute to streamline the development process of new drugs.

## Acknowledgments

We are grateful to Ms. Akiko Hisada and Mr. Hiroshi Sonoda for their helpful comments and suggestions. We also thank Ms. Kimie Mase and Ms. Yumi Mizunuma for technical assistance.

This study was funded by the in-house research budget of Hitachi High-Technologies Corp. and Hitachi, Ltd. The authors have no conflicts of interest.

## References

- [1] Ando H, Nishio Y, Ito K, Nakao A, Wang L, Zao YL, et al. Effect of endotoxin on P-glycoprotein-mediated biliary and renal excretion of Rhodamine-123 in rats. *Antimicrob Agents Chemother* 2001;45:3462–7.
- [2] Arrowsmith J, Miller P. Trial watch: phase II and phase III attrition rates 2011–2012. *Nat Rev Drug Discov* 2013;12:569.
- [3] Caron JM. Induction of albumin gene transcription in hepatocytes by extracellular matrix proteins. *Mol Cell Biol* 1990;10:1239–43.
- [4] Dunn JC, Yarmush ML, Koebe HG, Tompkins RG. Hepatocyte function and extracellular matrix geometry: long-term culture in a sandwich configuration. *FASEB J* 1989;3:174–7.
- [5] Emaus RK, Grunwald R, Lemasters JJ. Rhodamine 123 as a probe of transmembrane potential in isolated rat-liver mitochondria: spectral and metabolic properties. *Biochim Biophys Acta* 1986;850:436–48.
- [6] Fukuda H, Ohashi R, Tsuda-Tsukimoto M, Tamai I. Effect of plasma protein binding on in vitro-in vivo correlation of biliary excretion of drugs evaluated by sandwich-cultured rat hepatocytes. *Drug Metab Dispos* 2008;36:1275–82.
- [7] Ghibellini G, Vasist LS, Leslie EM, Heizer WD, Kowalsky RJ, Calvo BF, et al. In vitro-in vivo correlation of hepatobiliary drug clearance in humans. *Clin Pharmacol Ther* 2007;81:406–13.
- [8] Kremer BL, Staecher JL, Sawada N, Sattler GL, Hsia MT, Pitot HC. Use of a low-speed, iso-density percoll centrifugation method to increase the viability of isolated rat hepatocyte preparations. *In Vitro Cell Dev Biol* 1986;22:201–11.
- [9] LeCluyse EL, Audus KL, Hochman JH. Formation of extensive canalicular networks by rat hepatocytes cultured in collagen-sandwich configuration. *Am J Physiol* 1994;266:C1764–74.
- [10] Liu X, LeCluyse EL, Brouwer KR, Gan LS, Lemasters JJ, Stieger B, et al. Biliary excretion in primary rat hepatocytes cultured in a collagen-sandwich configuration. *Am J Physiol* 1999;277:G12–21.
- [11] Nakakariya M, Ono M, Amano N, Moriwaki T, Maeda K, Sugiyama Y. In vivo biliary clearance should be predicted by intrinsic biliary clearance in sandwich-cultured hepatocytes. *Drug Metab Dispos* 2012;40:602–9.
- [12] Nakanishi T, Shibue Y, Fukuyama Y, Yoshida K, Fukuda H, Shirasaka Y, et al. Quantitative time-lapse imaging-based analysis of drug-drug interaction mediated by hepatobiliary transporter, multidrug resistance-associated protein 2, in sandwich-cultured rat hepatocytes. *Drug Metab Dispos* 2011;39:984–91.
- [13] Paul SM, Mytelka DS, Dunwiddie CT, Persinger CC, Munos BH, Lindborg SR, et al. How to improve R&D productivity: the pharmaceutical industry's grand challenge. *Nat Rev Drug Discov* 2010;9:203–14.

- [14] Pfeifer ND, Yang K, Brouwer KL. Hepatic basolateral efflux contributes significantly to rosuvastatin disposition I: characterization of basolateral versus biliary clearance using a novel protocol in sandwich-cultured hepatocytes. *J Pharmacol Exp Ther* 2013;347:727–36.
- [15] Seglen PO. Preparation of isolated rat liver cells. *Methods Cell Biol* 1976;13: 29–83.
- [16] Takahashi R, Sonoda H, Tabata Y, Hisada A. Formation of hepatocyte spheroids with structural polarity and functional bile canaliculi using nanopillar sheets. *Tissue Eng Part A* 2010;16:1983–95.
- [17] Takahashi R, Hisada A, Sonoda H. Comprehensive expression analysis of primary hepatocytes under four different culture conditions. *Gene Expression Omnibus* accession number GSE38950. 2012.
- [18] Talamini MA, Kappus B, Hubbard A. Repolarization of hepatocytes in culture. *Hepatology* 1997;25:167–72.
- [19] Tian X, Zamek-Gliszczyński MJ, Zhang P, Brouwer KL. Modulation of multidrug resistance-associated protein 2 (Mrp2) and Mrp3 expression and function with small interfering RNA in sandwich-cultured rat hepatocytes. *Mol Pharmacol* 2004;66:1004–10.
- [20] Wu EY, Smith MT, Bellomo G, Di Monte D. Relationships between the mitochondrial transmembrane potential, ATP concentration, and cytotoxicity in isolated rat hepatocytes. *Arch Biochem Biophys* 1990;282:358–62.
- [21] Zamek-Gliszczyński MJ, Xiong H, Patel NJ, Turncliff RZ, Pollack GM, Brouwer KL. Pharmacokinetics of 5 (and 6)-carboxy-2',7'-dichlorofluorescein and its diacetate promoiety in the liver. *J Pharmacol Exp Ther* 2003;304:801–9.